



Unique Needs of the Adolescent

Elizabeth M. Alderman, MD, FSAHM, FAAP,^a Cora C. Breuner, MD, MPH, FAAP,^b COMMITTEE ON ADOLESCENCE

Adolescence is the transitional bridge between childhood and adulthood; it encompasses developmental milestones that are unique to this age group. Healthy cognitive, physical, sexual, and psychosocial development is both a right and a responsibility that must be guaranteed for all adolescents to successfully enter adulthood. There is consensus among national and international organizations that the unique needs of adolescents must be addressed and promoted to ensure the health of all adolescents. This policy statement outlines the special health challenges that adolescents face on their journey and transition to adulthood and provides recommendations for those who care for adolescents, their families, and the communities in which they live.

Adolescence, defined as 11 through 21 years of age,¹ is a critical period of development in a young person's life, one filled with distinctive and pivotal biological, cognitive, emotional, and social changes.² The World Health Organization³; the Office of Adolescent Health of the US Department of Health and Human Services⁴; the Health and Medicine Division of the National Academies of Sciences, Engineering, and Medicine (formerly the Institute of Medicine)^{5,6}; the *Lancet*,⁷ with 4 international academic institutions⁸; and the Society for Adolescent Health and Medicine⁹ have called for a closer examination of the unique health needs of adolescents. In 2018, *Nature* devoted an issue to the advances in the science of adolescence and called for ongoing further study of this important population.¹⁰ As a leader in adolescent health care, the American Academy of Pediatrics (AAP) is motivated to describe why adolescents are a unique and vulnerable population and why it is crucial that the AAP focus on adolescents' health concerns to optimize healthy development during the transition to adulthood. Addressing the unique needs of adolescents with disabilities is outside the scope of this statement; several statements specific to this population are available at <https://pediatrics.aappublications.org/collection/council-children-disabilities>. In addition, specific guidance around the transition to adult health care is not covered in this statement; please refer to the list of transition resources at the end of this document.

abstract

^aDivision of Adolescent Medicine, Department of Pediatrics, Albert Einstein College of Medicine and The Children's Hospital at Montefiore, Bronx, New York; and ^bDivision of Adolescent Medicine, Departments of Pediatrics and Orthopedics and Sports Medicine, University of Washington and Seattle Children's Hospital, Seattle, Washington

Policy statements from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, policy statements from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.

Drs Alderman and Breuner were equally responsible for writing and revising this policy statement with input from various internal and external reviewers as well as the Board of Directors, and both authors approved the final manuscript as submitted.

The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

DOI: <https://doi.org/10.1542/peds.2019-3150>

Address correspondence to Elizabeth M. Alderman, MD, FSAHM, FAAP. Email: ealderma@montefiore.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2019 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: No external funding.

To cite: Alderman EM, Breuner CC, AAP COMMITTEE ON ADOLESCENCE. Unique Needs of the Adolescent. *Pediatrics*. 2019;144(6):e20193150

The need for comprehensive health services for teenagers has been well documented since the 1990s.¹¹⁻¹³ The AAP advocates for the pediatrician to provide the medical home for adolescent primary care.¹⁴ Other professional societies, such as the Society for Adolescent Health and Medicine, the American Academy of Family Physicians, and the American College of Obstetricians and Gynecologists and school-based health initiatives (<https://www.sbh4all.org/>), recognize the unique needs of adolescents. These organizations recommend an increase in adolescent medicine training, along with the Accreditation Committee for Graduate Medical Education. The Accreditation Committee for Graduate Medical Education currently requires only 1 month of adolescent medicine training from a board-certified adolescent medicine specialist for all pediatric residency programs (adolescent medicine; [Core] IV.A.6.[b].[3].[a].[i]); there must be one educational unit).¹⁵ The importance of addressing the physical and mental health of adolescents has become more evident, with investigators in recent studies pointing to the fact that unmet health needs during adolescence and in the transition to adulthood predict not only poor health outcomes as adults but also lower quality of life in adulthood.¹⁶

HEALTH RISKS IN ADOLESCENCE

A hallmark of adolescence is a gradual development toward autonomy and individual adult decision-making. However, adolescents are often faced with situations for which they may not be prepared, and many are likely to be involved in risk-taking behaviors, such as use of alcohol, tobacco, and other drugs and engaging in unprotected sex. Most recently, there is increased concern about the rise in electronic cigarette use among

adolescence.¹⁷ In fact, most health care visits by adolescents to their pediatricians or other health care providers are to seek treatment of conditions or injuries that could have been prevented if screened for and addressed at an earlier comprehensive visit.¹⁸ Although some risk-taking behavior is considered normal in adolescence, engaging in certain types of risky behavior can have adverse and potentially long-term health consequences. The majority of mortality and morbidity during adolescence, which can be prevented, is attributable to unintentional injuries, suicide, and homicide.¹⁹ Approximately 72% of deaths among adolescents are attributable to injuries from motor vehicle crashes, other unintentional and intentional injuries, injuries caused by firearms, injuries influenced by use of alcohol and illicit substances, homicide, or suicide.^{20,21} These causes of death greatly surpass medical etiologies such as cancer, HIV infection, and heart disease in the United States and other industrialized nations.²¹

The AAP *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents* recommends a strength-based approach to screening and counseling around these behaviors that lead to mortality and morbidity in adolescents.^{1,22} However, according to the National Ambulatory Medical Care Survey and the National Hospital Ambulatory Medical Care Survey, only 39% of adolescents received any type of preventive counseling during ambulatory visits.²³ Seventy-one percent of teenagers reported at least 1 potential health risk, yet only 37% of these teenagers reported discussing any of these risks with their pediatrician or primary care physician. Clearly, screening for and counseling around these high-risk behaviors needs to be improved.²⁴

New screening codes for depression, substance use, and alcohol and tobacco use as well as brief intervention services may provide opportunities to receive payment for the services pediatricians are providing to adolescents. These include 96127, brief emotional and behavioral assessment (eg, depression inventory, attention-deficit/hyperactivity disorder scale) with scoring and documentation, per standardized instrument, and 96150, health and behavior assessment (eg, health-focused clinical interview, behavioral observations, psychophysiological monitoring, health-oriented questionnaires).²⁵ However, it is important to recognize that coding for specific diagnoses may be challenging if the patient does not want his or her parent(s) to know the reasons for the clinical visit. Adolescent visits and documentation of visits are confidential to promote better access and to protect the rights of adolescents.²⁶

Another trend in the health status of adolescents (reflecting technological advances in pediatric medical care) is the increasing number of pediatric patients with chronic medical conditions and developmental challenges who enter adolescence. Adolescents with chronic conditions face developmental challenges similar to their healthy peers but may have special educational, vocational, and transitional concerns because of their medical issues.²⁷

The prevalence of chronic medical conditions and developmental and physical disabilities in adolescents is difficult to assess because of the variation of study methodologies and categorical versus noncategorical approaches to the epidemiology of chronic illness.²⁸ According to the National Survey of Children's Health, funded by the US Department of Health and Human Services, almost 31% of adolescents have 1 moderate to severe chronic illness, such as asthma or a mental health

condition.²⁹ Other common chronic illnesses include obesity, cancer, cardiac disease, HIV infection, spastic quadriplegia, and developmental disabilities.³⁰⁻³² One in 4 adolescents with chronic illness has at least 1 unmet health need that may affect physical growth and development, including puberty and overall health status as well as future adult health.³³

Within pediatric practice, integrating adolescent-centered, family-involved approaches into the care of adolescents as well as culturally competent and effective approaches (as outlined in *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*) has the potential not only to identify threats to well-being but also to create a space to work with families to bolster opportunities for optimal development of all children.¹ When considering the health challenges adolescents face, it is imperative to take into account not only the ethnic and racial diversity of the adolescent population in the United States but also the social and ecologic factors (eg, socioeconomic status, family composition, parental education and engagement, neighborhood and school environment, religion, earlier childhood trauma and toxic stress, and access to health care).

The Search Foundation has conducted research that suggests that for minority youth, a positive ethnic identity is a critical spark for emergence of the required developmental assets to enable adolescents to develop into successful and contributing adults.^{34,35} This theory is supported by a recent study in *The Journal of Pediatrics* that suggests minority youth are still prone to depression because of isolation and discrimination faced during adolescence while navigating neighborhood and school environments, even when they have educated and supportive parents.³⁶

African American male adolescents have the highest rates of mortality, followed by American Indian, white, Hispanic, and Asian American or Pacific Islander male adolescents, pointing to racial and ethnic disparities in adolescent health and the potential to achieve a healthy adulthood.³⁷

The AAP has previously published policy statements addressing the unique strengths and health disparities that exist for specific groups of adolescents, such as lesbian, gay, bisexual, and transgender youth and those in the juvenile justice system, foster care, and the military.³⁸⁻⁴² Pediatricians must pay attention to how care is delivered to the increasingly diverse adolescent populations to prevent a decline in health status and increase in health care disparities.

UNIQUE BIOLOGICAL AND PSYCHOSOCIAL CHANGES OCCURRING DURING ADOLESCENCE

Biological and psychosocial changes that occur during adolescence make this age group unique. Research describing the timing and physiology of puberty has been invaluable in revealing not only differences between racial groups but also between adolescents with different chronic conditions.⁴³⁻⁴⁶

Puberty

Puberty is the hallmark of physiologic progression from child to adult body habitus. Chronic conditions, such as obesity and intracranial lesions, or trauma may cause early puberty, which may put the adolescent at risk for engagement in higher-risk behaviors at an earlier age.⁴⁴ Delayed puberty is often a variant of normal development but may also be seen in adolescents with inflammatory bowel disease, eating disorders, and chronic conditions that create malnutrition as well as adolescents who have undergone treatment of malignancies.

Comorbid mental disease (eg, an eating disorder that causes delayed puberty) or medication for psychiatric illness that causes obesity, which may cause early puberty, can complicate optimal adolescent psychosocial development.

Brain Development

The work of Giedd⁴⁷ and others shows that brain development during adolescence is ongoing and affects behavior and health. Because of changes in signaling that relate to the reward system in which the brain motivates behavior and the continuing maturation of the parts of the brain that regulate impulse control, adolescents may have a propensity to be involved with high-risk behaviors and have heightened response to emotionally loaded situations. In addition, adverse childhood experiences can have an impact on brain development, affecting behaviors and health during adolescence.⁴⁸ During adolescence, there is a “pruning” of gray matter and synapses, which makes the brain more efficient.⁴⁷ White matter increases throughout adolescence, which allows the older adolescent and adult brain to conduct more-complex cognitive tasks and adaptive behavior.⁴⁹

Increasingly, studies show that the adolescent brain responds to alcohol and illicit substances differently than adults.^{50,51} This difference may explain the increased risk of binge drinking as well as greater untoward cognitive effects of alcohol and marijuana.

Sexual Orientation and Gender Identification

Sexual (and gender) development is a process that starts early in childhood and involves negotiating and experimenting with identity, relationships, and roles. In early adolescence, people begin to

recognize or become aware of their sexual orientation.^{52,53}

However, some adolescents are still unsure of their sexual attractions, and others struggle with their known sexual attraction. Adolescence is a time of identity formation and experimentation, so labels that one uses for their sexual orientation (eg, gay, straight, bisexual, etc) often do not correlate to actual sexual behaviors and partners. Sexual orientation and behaviors should be assessed by the pediatrician without making assumptions. Adolescents should be allowed to apply and explain the labels they choose to use for sexuality and gender using open-ended questions.^{54–56}

Sexual minority adolescents may engage in heterosexual practices, and heterosexual adolescents may engage in same-sex sexual activity. Depending on their specific behaviors and the gender of various partners, all sexually active adolescents may be at risk for sexually transmitted infections and unplanned pregnancy. Sexual minority youth are at higher risk of sexually transmitted infections and unplanned pregnancy, often because they do not receive education that applies to their sexual behaviors and are less likely to be screened appropriately (<http://www.cdc.gov/healthyyouth/disparities/smy.htm>).^{57,58}

Sexual minority and transgender youth, because of the stigma they face, are also at higher risk of mental health problems, including depression and suicidality, altered body image, and substance use.³⁸

There is strong evidence that when sexual minority and transgender youth feel they cannot express their true selves, they go underground by either hiding or denying their attractions and identity.⁵⁹ When this is combined with reinforcing parental rejection, bullying, etc, it is believed to lead to internalization, low self-

esteem, and ultimately, depression and suicide.⁵⁹ Using an explanation like this places the problem on the societal context, not the adolescent or his or her identity.^{38,39,60,61}

A relatively higher proportion of homeless adolescents are lesbian, gay, bisexual, transgender, and queer or questioning youth.⁶¹ They leave their family homes because of abuse or having been thrown out. These adolescents are at high risk for victimization and often need to engage in unsafe sexual practices to provide themselves food and shelter.⁶¹

Mental health problems may become more pronounced when sexual minority teenagers come out during adolescence to unsupportive family members and friends or health care providers.³⁸ These youth are more likely to experience violence both in their homes and in their schools and communities. Studies have shown that sexual minority youth reveal higher rates of tobacco, alcohol, marijuana, and other illicit substance use.⁶²

Most adolescents identify by and express a gender that conforms to their anatomic sex. However, some adolescents experience gender dysphoria with their anatomic sex when entering puberty. As they consider transgender options, they are at an increased risk of mental or emotional health problems, including depression and suicidality, victimization and violence, eating disorders, substance use, and unaccepting or intolerant family members and peers. Crucial to the successful navigation of gender dysphoria issues are health care providers who can assist transgender youth and families to achieve safe, healthy transitioning both in the postponement of puberty, when indicated, and in transitioning to preferred gender with psychosocial and behavioral support.⁵⁹

Legal Status

Adolescence heralds a change of legal status, in which the age of 18 or 19 years transforms legal status from minors to adults with full legal privileges and obligations related to health care. However, certain states afford minors the right to confidentiality and consent to or for reproductive and mental health and substance use treatment confidential health services.^{26,63} Generally, minors may receive confidential screening and care for sexually transmitted infections in all 50 states and the District of Columbia. However, accessing contraception to prevent unwanted pregnancy as well as the ability to self-consent to pregnancy options counseling, prenatal care, and termination of pregnancy vary between states.⁶⁴ These discrepancies also exist in accessing outpatient mental health and substance use services. Many adolescents in need of these services do not know they may have the right to access them on their own and may avoid interaction with the health care system to assist with reproductive and mental health concerns.¹⁶ Delaying such care leads to adverse health outcomes.¹⁶ A recent survey confirms that adolescents value private time with their health care providers, with confidentiality assurances by health care providers.⁶⁵ The need for office policies in negotiating private time was suggested. Moreover, health care providers reported needing more education in the provision of confidential services.⁶⁶ Adolescents in foster care may also be limited in their autonomous access to confidential services, which varies state to state.⁴¹ In certain states, pregnant and parenting adolescents may have the right to consent for their care and the care of their child (<https://www.guttmacher.org/state-policy/explore/minors-rights-parents>, [4](https://www.schoolhouseconnection.org/state-laws-on-minor-consent-for-routine-</p></div><div data-bbox=)

medical-care/). Few adolescents are considered emancipated minors and, thereby, entitled to all legal privileges of adults.⁶⁷

Mental Health and Emotional Well-being

Mental health and emotional well-being, in combination with issues pertaining to sexual and reproductive health, violence and unintentional injury, substance use, eating disorders, and obesity, create potential challenges to adolescents' healthy emotional and physical development.⁶⁸ Approximately 20% of adolescents have a diagnosable mental health disorder.⁶⁹ Many mental health disorders present initially during adolescence. Twenty-five percent of adults with mood disorder had their first major depressive episode during adolescence.⁷⁰

Suicide is the second leading cause of death in adolescents, resulting in more than 5700 deaths in 2016.⁷¹ Between 2007 and 2016, the overall suicide rate for children and adolescents ages 10 to 19 years increased by 56%.⁷¹ Older adolescents (15–19 years of age) are at an increased risk of suicide, with a rate of 5 in 100 000 for girls and 20 in 100 000 for boys.⁷¹ According to the 2017 Youth Risk Behavior Survey of high school students, 7.4% of high school students attempted suicide in the last 12 months, and 13.6% made a suicide plan.⁷² Adolescents with parents in the military were at increased risk of suicidal ideation (odds ratio [OR]: 1.43; 95% confidence interval [CI]: 1.37–1.49), making a plan to harm themselves (OR: 1.19; 95% CI: 1.06–1.34), attempting suicide (OR: 1.67; 95% CI: 1.43–1.95), and an attempted suicide that required medical treatment.⁷³

Eating disorders typically present in the adolescent years. Although the incidence of eating disorders is low compared with depression, anxiety,

and other mental health problems, these problems are often comorbid with eating disorders.⁷⁴ Moreover, the incidence of anorexia nervosa, bulimia nervosa, and other disordered eating is becoming more prevalent in formerly obese teenagers, male teenagers, and teenagers from lower socioeconomic groups.^{75–77}

Teenagers with mental health issues may have subsequent poor school performance, school dropout, difficult family relationships, involvement in the juvenile justice system, substance use, and high-risk sexual behaviors.⁷⁸ Almost 70% of youth in the juvenile justice system have a diagnosed mental health disorder.^{79,80}

Rates of serious mental health disorders among homeless youth range from 19% to 50%.^{81,82} Homeless youth have a high need for treatment but rarely use formal treatment programs for medical, mental, and substance use services.⁸¹ Confidentiality is also an issue for adolescents, as evidenced by the fact that in adolescents to whom confidentiality is not assured, there is a higher prevalence of depressive symptoms, suicidal thoughts, and suicide attempts.⁸³ There is a paucity of adequately trained mental health professionals to care for adolescents with these mental health challenges.⁸⁴ In addition, coverage for mental health services by insurance plans can be variable.⁷⁸

Morbidity From High-Risk Sexual Activity

Multiple factors, including the increase in use of long-acting reversible contraception, have resulted in the teenage pregnancy rate decreasing in the United States over the past 20 years.^{85,86} However, pregnancy still contributes to delays in educational and career success for adolescents. Moreover, pregnant teenagers are more likely to delay

seeking medical care, putting them at risk for pregnancy-related health problems and putting their children at risk for prematurity and other negative birth outcomes.⁸⁷

Adolescents continue to have the highest rates of sexually transmitted infections (eg, gonorrhea and *Chlamydia*).⁸⁸ Although screening most sexually active adolescents for *Chlamydia* infection is covered by the Patient Protection and Affordable Care Act (Pub L No. 111–148 [2010]) and recommended by the AAP *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, adolescent concerns about billing and confidentiality are obstacles to medical screening.^{1,89} Pediatricians can refer to AAP guidance to find appropriate codes for payment for providing adolescent health services (https://www.aap.org/en-us/Documents/coding_factsheet_adolescenthealth.pdf).

THE ADOLESCENT MEDICAL HOME

Consideration of the unique health risks as well as the biological and psychosocial elements of adolescence allows the AAP-endorsed patient-centered medical home (PCMH) to serve as an ideal conceptual framework by which a primary care practice can maximize the quality, efficiency, and patient experience of care. In 2007, the AAP joined the American Academy of Family Physicians, the American College of Physicians, and the American Osteopathic Association to endorse the “Joint Principals of the Patient-Centered Medical Home,” which describes 7 core characteristics: (1) personal physician for every patient; (2) physician-directed medical practice; (3) whole person orientation; (4) care is coordinated and/or integrated; (5) quality and safety are hallmarks of PCMH care; (6)

enhanced access to care; and (7) appropriate payment for providing PCMH care.⁹⁰ The AAP, American Academy of Family Physicians, and American College of Physicians assert that optimal health care is achieved when each person, at every age, receives developmentally appropriate care.⁹¹ Pediatricians provide quality adolescent care when they maintain relationships with families and with their patients and, thus, help patients develop autonomy, responsibility, and an adult identity.⁹² Issues unique to adolescence to consider within the PCMH model include the following: adolescent-oriented developmentally appropriate care, which may require longer appointment times; confidentiality of health care visits, health records, billing, and the location where adolescents receive care; providers who offer such care; and the transition to adult care.^{91,93} Moreover, using a strengths-based approach in the care of adolescents, as well as capitalizing on resiliency, is instrumental to maintaining the health of the individual adolescent.⁹⁴

Schools have an important role for adolescents who either do not have access to a PCMH or do not use their access to receive recommended preventive services. School-based health centers and school-based mental health services can meet the needs of adolescents who do not have a PCMH or can coordinate school-based health services with the PCMH if the student has one. School nurses can help identify and refer adolescents who need these services.⁹⁵

Financing health care of the adolescent can be challenging. Please see the detailed AAP policy statement on reforms in health care financing with the ultimate goal to improve the health care of all adolescents.⁹²

RECOMMENDATIONS

On the basis of the unique biological and psychosocial aspects of adolescence, the AAP supports the following:

1. continued recognition by international and national organizations, including the AAP, the Society for Adolescent Health and Medicine, the American College of Obstetricians and Gynecologists, the North American Society for Pediatric and Adolescent Gynecology, and the American Academy of Family Physicians, of the need for policies and advocacy related to adolescent health and well-being;
2. sustained funding for research to further elucidate the biological basis of the growth and development of adolescents and how they affect adolescent behavior;
3. educational programs and adequate financial compensation for pediatricians and other health care professionals to support them in providing evidence-based, quality primary care for adolescents;
4. pediatricians receiving training on how to maintain the clinical setting as a “safe space,” particularly in terms of confidentiality, especially when working with lesbian, gay, bisexual, transgender, and queer or questioning adolescents;
5. the role of schools, including school nurses and school-based health centers, and their role in promoting healthy adolescent development and providing access to health care;
6. further education, training, and advocacy for mental health care services that specifically address the needs of adolescents, preferably as part of a medical home model, stressing the

importance of mental health for all youth;

7. federal confidentiality protection for mental health and reproductive services, as is currently provided in many states;
8. innovative postresidency training programs to increase the number of adolescent-trained pediatric providers in the workforce;
9. improved access to medical homes for all adolescents to ensure access to preventive medical care;
10. affiliation of middle and high schools with a physician trained to care for adolescents, unless the student already has access to comprehensive adolescent health services;
11. education for pediatricians so that they are aware of the laws regarding confidential care of adolescents in their states; and
12. familiarity with community resources for confidential reproductive and mental health care if they cannot provide confidential care themselves. Pediatricians who are unable to provide these services should learn about local community resources that provide confidential reproductive and mental health care.

The AAP recommends the following strategies targeted at improving financing for the health care of adolescents:

1. Federal and state agencies should increase their efforts to further reduce the number of adolescents who are not insured or who lack comprehensive and affordable health insurance.
2. The Centers for Medicare and Medicaid Services should implement its regulatory authority to update its standards for essential health benefits, as defined in the Patient Protection

and Affordable Care Act, in the 2 categories of mental and behavioral health services and pediatric services. These essential health benefits should be consistent with the full scope of benefits outlined in *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents* (including health supervision visits, recommended immunizations, screening for high-risk conditions, and adequate counseling and treatment of conditions related to sexual, reproductive, mental, and behavioral health and substance use disorder). In this way, all adolescents can access the full range of services needed during this developmentally critical period to secure optimal physical and mental health on entry into midadulthood.

3. All health plans should provide preventive services without member cost sharing. In addition, to reduce financial barriers to care for adolescents, payers should limit the burden on families by reducing or eliminating copayments and eliminating coinsurance for visits related to anticipatory guidance and/or treatment of sexual and reproductive health, behavioral health, and immunization visits.
4. To provide sufficient payment to physicians and other health care providers for medical services to adolescents, insurers' claims systems should recognize and pay for all preventive medicine *Current Procedural Terminology* codes related to services for health and behavior assessment, counseling, risk screening, and/or appropriate interventions recommended in *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. These services should not be bundled under a single health maintenance

Current Procedural Terminology code.

5. Government and private insurance payers should increase the relative value unit allocation and level of payment for pediatricians delivering care and clinical preventive services to adolescents to a level that is commensurate with the time and effort expended, including health maintenance services, screening, and counseling.
6. The Centers for Medicare and Medicaid Services should mandate that payers provide enhanced access to cost-effective and clinically sound behavioral health services for adolescents, ensure that payment for all mental health services is more equitable with payment provided for medical and surgical services, and ensure that pediatricians are paid for mental health services provided during health maintenance and follow-up visits.

APPENDIX: ONLINE RESOURCES

AMERICAN ACADEMY OF PEDIATRICS

- Confidentiality Protections for Adolescents and Young Adults in the Health Care Billing and Insurance Claims Process: <http://pediatrics.aappublications.org/content/137/5/e20160593>
- Office-Based Care for Lesbian, Gay, Bisexual, Transgender, and Questioning Youth: <http://pediatrics.aappublications.org/content/pediatrics/132/1/e297.full.pdf>

SOCIETY FOR ADOLESCENT HEALTH AND MEDICINE

Resources for adolescents and parents are online resources aimed specifically at adolescents and their parents. Health care providers and youth-serving professionals can offer these additional resources or print a 1-page reference sheet (PDF) for adolescents and parents looking for

additional information, including support groups, peer networks, helplines, treatment locators, and advocacy opportunities.

- Mental Health Resources for Adolescents: <http://www.adolescenthealth.org/Topics-in-Adolescent-Health/Mental-Health/Mental-Health-Resources-For-Adolesc.aspx>
- Mental Health Resources for Parents of Adolescents: <http://www.adolescenthealth.org/Topics-in-Adolescent-Health/Mental-Health/Mental-Health-Resources-For-Parents-of-Adolescents.aspx>
- Substance Use Resources for Adolescents: <http://www.adolescenthealth.org/Topics-in-Adolescent-Health/Substance-Use/Substance-Use-Resources-For-Adolesc.aspx>
- Substance Use Resources for Parents of Adolescents: <http://www.adolescenthealth.org/Topics-in-Adolescent-Health/Substance-Use/Substance-Use-Resources-For-Parents-of-Adolesc.aspx>
- Confidentiality in Health Care Resources for Adolescents and Parents of Adolescents: <http://www.adolescenthealth.org/Topics-in-Adolescent-Health/Confidentiality/Confidentiality-Resources-For-Adolesc.aspx>
- Sexual and Reproductive Health Resources for Adolescents: <http://www.adolescenthealth.org/Topics-in-Adolescent-Health/Sexual-Reproductive-Health/Sexual-Reproductive-Health-Resources-For-Adolesc.aspx>
- Sexual and Reproductive Health Resources for Parents of Adolescents: <http://www.adolescenthealth.org/Topics-in-Adolescent-Health/Sexual-Reproductive-Health/SandRH-Resources-For-Parents-of-Adolesc.aspx>
- Physical and Psychosocial Development Resources for

Adolescents: <http://www.adolescenthealth.org/Topics-in-Adolescent-Health/Physical-and-Psychosocial-Development/Physical-Psychosocial-Development-Resources-For-Adolesc.aspx>

- Physical and Psychosocial Development Resources for Parents of Adolescents: <http://www.adolescenthealth.org/Topics-in-Adolescent-Health/Physical-and-Psychosocial-Development/Physical-Psych-Resources-For-Parents-of-Adolesc.aspx>

TRANSITION RESOURCES

General Resources

- National Health Care Transition Center (www.gottransition.org)
- Family Voices, Inc (www.familyvoices.org)
- National Alliance to Advance Adolescent Health (www.thenationalalliance.org)

Transition Care Plans

- AAP/National Center for Medical Home Implementation (www.medicalhomeinfo.org/how/care_delivery/transitions.aspx)
- British Columbia Ministry of Children and Family Development, "Transition Planning for Youth With Special Needs" (www.mcf.gov).

bc.ca/spec_needs/pdf/support_guide.pdf

- University of Washington, Adolescent Health Transition Project (<http://depts.washington.edu/healthtr>)

Transition Assessment and Evaluation Tools

- AAP/National Center for Medical Home Implementation (www.medicalhomeinfo.org/health/trans.html)
- JaxHATS, evaluation tools for youth and caregivers and training materials for medical providers (www.jaxhats.ufl.edu/docs)
- Texas Children's Hospital transition template (<http://leah.mchtraining.net/bcm/resources/tracs>)
- Carolina Health and Transition Project (www.mahec.net/quality/chat.aspx?a=10)
- University of Washington, Adolescent Health Transition Project (<http://depts.washington.edu/healthtr>)
- Wisconsin Community of Practice on Transition (www.waisman.wisc.edu/wrc/pdf/pubs/THCL.pdf)

NATIONAL ALLIANCE TO END HOMELESSNESS

- <http://www.endhomelessness.org/>

LEAD AUTHORS

Elizabeth M. Alderman, MD, FSAHM, FAAP
Cora C. Breuner, MD, MPH, FAAP

COMMITTEE ON ADOLESCENCE, 2017–2018

Cora Breuner, MD, MPH, FAAP, Chairperson
Elizabeth M. Alderman, MD, FSAHM, FAAP
Laura K. Grubb, MD, MPH, FAAP
Makia E. Powers, MD, MPH, FAAP
Krishna Upadhy, MD, FAAP
Stephenie B. Wallace, MD, FAAP

LIAISONS

Laurie Hornberger, MD, MPH, FAAP – *Section on Adolescent Health*
Liwei L. Hua, MD, PhD – *American Academy of Child and Adolescent Psychiatry*
Margo A. Lane, MD, FRCPC, FAAP – *Canadian Paediatric Society*
Meredith Loveless, MD, FACOG – *American College of Obstetricians and Gynecologists*
Seema Menon, MD – *North American Society of Pediatric and Adolescent Gynecology*
CDR Lauren B. Zapata, PhD, MSPH – *Centers for Disease Control and Prevention*

STAFF

Karen S. Smith
James D. Baumberger, MPP

ABBREVIATIONS

AAP: American Academy of Pediatrics
CI: confidence interval
OR: odds ratio
PCMH: Patient-Centered Medical Home

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

REFERENCES

1. Hagan JF, Shaw JS, Duncan PM, eds. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 4th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2017
2. Shlafer R, Hergenroeder AC, Jean Emans S, et al. Adolescence as a critical stage in the MCH Life Course Model: commentary for the Leadership Education in Adolescent Health (LEAH) interdisciplinary training program projects. *Matern Child Health J*. 2014; 18(2):462–466
3. World Health Organization. *Health for the World's Adolescents: A Second Chance for a Second Decade*. Geneva, Switzerland: World Health Organization; 2014. Available at: www.who.int/maternal_child_adolescent/documents/second-decade/en/. Accessed March 27, 2017
4. United States Department of Health and Human Services. About the office of adolescent health. Available at: www.hhs.gov/ash/oah/about-us/index.html. Accessed March 27, 2017
5. Institute of Medicine. *Child and Adolescent Health and Health Care Quality: Measuring What Matters*. Washington, DC: National Academies Press; 2011. Available at: <http://nationalacademies.org/hmd/reports/2011/child-and-adolescent-health-and-health-care-quality.aspx>. Accessed March 27, 2017

6. Institute of Medicine. *Adolescent Health Services: Missing Opportunities*. Washington, DC: National Academies Press; 2008. Available at: <http://nationalacademies.org/hmd/reports/2008/adolescent-health-services-missing-opportunities.aspx>. Accessed March 27, 2017
7. Patton GC, Sawyer SM, Santelli JS, et al. Our future: a *Lancet* commission on adolescent health and wellbeing. *Lancet*. 2016;387(10036):2423–2478
8. Patton GC, Ross DA, Santelli JS, et al. Next steps for adolescent health: a *Lancet* Commission. *Lancet*. 2014; 383(9915):385–386
9. English A, Park MJ, Shafer MA, Kreipe RE, D'Angelo LJ. Health care reform and adolescents-an agenda for the lifespan: a position paper of the Society for Adolescent Medicine. *J Adolesc Health*. 2009;45(3):310–315
10. Special issue: adolescence. *Nature*. 2018;554(7693):403–561
11. Elster AB. Comparison of recommendations for adolescent clinical preventive services developed by national organizations. *Arch Pediatr Adolesc Med*. 1998;152(2): 193–198
12. Ressel G; American Academy of Family Physicians. Introduction to AAFP summary of recommendations for periodic health examinations. *Am Fam Physician*. 2002;65(7):1467
13. Blum RW, Bastos FI, Kabiru CW, Le LC. Adolescent health in the 21st century. *Lancet*. 2012;379(9826):1567–1568
14. American Academy of Pediatrics Ad Hoc Task Force on Definition of the Medical Home: the medical home. *Pediatrics*. 1992;90(5):774
15. Accreditation Council on Graduate Medical Education. ACGME program requirements for graduate medical education in pediatrics. 2013. Available at: <https://www.acgme.org/Specialties/Program-Requirements-and-FAQs-and-Applications/pfcatid/16/Pediatrics>. Accessed March 27, 2017
16. Hargreaves DS, Elliott MN, Viner RM, Richmond TK, Schuster MA. Unmet health care need in US adolescents and adult health outcomes. *Pediatrics*. 2015; 136(3):513–520
17. Jenssen BP, Walley SC; Section on Tobacco Control. E-cigarettes and similar devices. *Pediatrics*. 2019;143(2): e20183652
18. Nordin JD, Solberg LI, Parker ED. Adolescent primary care visit patterns. *Ann Fam Med*. 2010;8(6):511–516
19. Heron M; Centers for Disease Control and Prevention. Deaths: leading causes for 2013. *Natl Vital Stat Rep*. 2016;65(2): 1–95
20. Centers for Disease Control and Prevention. National Center for Health Statistics Compressed mortality rate 1999-2010. 2012. Available at: <http://wonder.cdc.gov/ucd-icd10.html>. Accessed March 27, 2017
21. Murphy SL, Xu J, Kochanek KD, Curtin SC, Arias E. Deaths: final data for 2015. *Natl Vital Stat Rep*. 2017;66(6): 1–75
22. American Academy of Pediatrics. Bright futures. Adolescence Tools. Available at: <https://brightfutures.aap.org/materials-and-tools/tool-and-resource-kit/Pages/adolescence-tools.aspx>. Accessed March 27, 2017
23. Ma J, Wang Y, Stafford RSUS. U.S. adolescents receive suboptimal preventive counseling during ambulatory care. *J Adolesc Health*. 2005;36(5):441
24. Chung PJ, Lee TC, Morrison JL, Schuster MA. Preventive care for children in the United States: quality and barriers. *Annu Rev Public Health*. 2006;27: 491–515
25. American Academy of Pediatrics. *Coding for Pediatrics 2016: A Manual for Pediatric Documentation and Payment*. Elk Grove Village, IL: American Academy of Pediatrics; 2016
26. Society for Adolescent Health and Medicine; American Academy of Pediatrics. Confidentiality protections for adolescents and young adults in the health care billing and insurance claims process. *J Adolesc Health*. 2016; 58(3):374–377
27. Hale DR, Bevilacqua L, Viner RM. Adolescent health and adult education and employment: a systematic review. *Pediatrics*. 2015;136(1):128–140
28. Michaud PA, Suris JC, Viner R. *The Adolescent with a Chronic Condition: Epidemiology, Developmental Issues and Health Care Provision*. Geneva, Switzerland: World Health Organization; 2007. Available at http://apps.who.int/iris/bitstream/10665/43775/1/9789241595704_eng.pdf. Accessed March 27, 2017
29. Child and Adolescent Health Measurement Initiative. Data Resource Center for Child and Adolescent Health: 2016 National Survey of Children's Health data query. Available at: www.childhealthdata.org. Accessed March 27, 2017
30. Newacheck PW, McManus MA, Fox HB. Prevalence and impact of chronic illness among adolescents. *Am J Dis Child*. 1991;145(12):1367–1373
31. Denny S, de Silva M, Fleming T, et al. The prevalence of chronic health conditions impacting on daily functioning and the association with emotional well-being among a national sample of high school students. *J Adolesc Health*. 2014; 54(4):410–415
32. Compas BE, Jaser SS, Dunn MJ, Rodriguez EM. Coping with chronic illness in childhood and adolescence. *Annu Rev Clin Psychol*. 2012;8: 455–480
33. Child and Adolescent Health Measurement Initiative, Data Resource Center for Child and Adolescent Health. *National Survey of Children's Health (2016–2017)*. Portland, OR: Child and Adolescent Health Measurement Initiative, the Data Resource Center for Child and Adolescent Health; 2009–2010. Available at: <http://childhealthdata.org/browse/survey/results?q=1624&r=1&g=86>. Accessed March 27, 2017
34. Search Foundation. Relationships that matter to America's teens. Available at: www.search-institute.org/sites/default/files/a/TeenVoice2010.pdf. Accessed March 27, 2017
35. Search Institute. Developmental assets. Available at: www.search-institute.org/research/developmental-assets. Accessed March 27, 2017
36. Cheng ER, Cohen A, Goodman E. The role of perceived discrimination during

- childhood and adolescence in understanding racial and socioeconomic influences on depression in young adulthood. *J Pediatr*. 2015;166(2):370–377.e1
37. Centers for Disease Control and Prevention, National Center for Health Statistics. Vital statistics online data portal. Available at: www.cdc.gov/nchs/data_access/Vitalstatsonline.htm. Accessed March 27, 2017
 38. Committee On Adolescence. Office-based care for lesbian, gay, bisexual, transgender, and questioning youth. *Pediatrics*. 2013;132(1):198–203
 39. Levine DA; Committee On Adolescence. Office-based care for lesbian, gay, bisexual, transgender, and questioning youth. *Pediatrics*. 2013;132(1). Available at: www.pediatrics.org/cgi/content/full/132/1/e297
 40. Committee on Adolescence. Health care for youth in the juvenile justice system. *Pediatrics*. 2011;128(6):1219–1235
 41. Council on Foster Care, Adoption, and Kinship Care; Committee on Adolescence; Council on Early Childhood. Health care issues for children and adolescents in foster care and kinship care. *Pediatrics*. 2015;136(4). Available at: www.pediatrics.org/cgi/content/full/136/4/e1131
 42. Siegel BS, Davis BE; Committee on Psychosocial Aspects of Child and Family Health and Section on Uniformed Services. Health and mental health needs of children in US military families. *Pediatrics*. 2013;131(6). Available at: www.pediatrics.org/cgi/content/full/131/6/e2002
 43. Herman-Giddens ME, Slora EJ, Wasserman RC, et al. Secondary sexual characteristics and menses in young girls seen in office practice: a study from the Pediatric Research in Office Settings network. *Pediatrics*. 1997;99(4):505–512
 44. Suris JC, Michaud PA, Viner R. The adolescent with a chronic condition. Part I: developmental issues. *Arch Dis Child*. 2004;89(10):938–942
 45. Romer D, Reyna VF, Satterthwaite TD. Beyond stereotypes of adolescent risk taking: placing the adolescent brain in developmental context. *Dev Cogn Neurosci*. 2017;27:19–34
 46. Suleiman AB, Dahl RE. Leveraging neuroscience to inform adolescent health: the need for an innovative transdisciplinary developmental science of adolescence. *J Adolesc Health*. 2017;60(3):240–248
 47. Giedd JN. Structural magnetic resonance imaging of the adolescent brain. *Ann N Y Acad Sci*. 2004;1021(1):77–85
 48. Hughes K, Bellis MA, Hardcastle KA, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *Lancet Public Health*. 2017;2(8):e356–e366
 49. Giedd JN. The teen brain: insights from neuroimaging. *J Adolesc Health*. 2008;42(4):335–343
 50. Bava S, Tapert SF. Adolescent brain development and the risk for alcohol and other drug problems. *Neuropsychol Rev*. 2010;20(4):398–413
 51. Spear LP. The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev*. 2000;24(4):417–463
 52. Frankowski BL; American Academy of Pediatrics Committee on Adolescence. Sexual orientation and adolescents. *Pediatrics*. 2004;113(6):1827–1832
 53. Spigarelli MG. Adolescent sexual orientation. *Adolesc Med State Art Rev*. 2007;18(3):508–518, vii
 54. Zimmerman MA. Resiliency theory: a strengths-based approach to research and practice for adolescent health. *Health Educ Behav*. 2013;40(4):381–383
 55. Duncan PM, Garcia AC, Frankowski BL, et al. Inspiring healthy adolescent choices: a rationale for and guide to strength promotion in primary care. *J Adolesc Health*. 2007;41(6):525–535
 56. Benson PL, Scales PC, Hamilton SF, Sesma A Jr. Positive Youth Development: Theory, Research and Applications. In: Damon W, Lerner R, eds. *Handbook of Child Psychology*. Hoboken, NJ: John Wiley & Sons; 2007:894–941
 57. Igartua K, Thombs BD, Burgos G, Montoro R. Concordance and discrepancy in sexual identity, attraction, and behavior among adolescents. *J Adolesc Health*. 2009;45(6):602–608
 58. Goodenow C, Szalacha LA, Robin LE, Westheimer K. Dimensions of sexual orientation and HIV-related risk among adolescent females: evidence from a statewide survey. *Am J Public Health*. 2008;98(6):1051–1058
 59. Rafferty J; Committee on Psychosocial Aspects of Child and Family Health; Committee on Adolescence; Section on Lesbian, Gay, Bisexual, and Transgender Health and Wellness. Ensuring comprehensive care and support for transgender and gender-diverse children and adolescents. *Pediatrics*. 2018;142(4):e20182162
 60. Kitts RL. Gay adolescents and suicide: understanding the association. *Adolescence*. 2005;40(159):621–628
 61. Keuroghlian AS, Shtasel D, Bassuk EL. Out on the street: a public health and policy agenda for lesbian, gay, bisexual, and transgender youth who are homeless. *Am J Orthopsychiatry*. 2014;84(1):66–72
 62. Parsons JT, Kelly BC, Weiser JD. Initiation into methamphetamine use for young gay and bisexual men. *Drug Alcohol Depend*. 2007;90(2–3):135–144
 63. Alan Guttmacher Institute. An overview of consent to reproductive health services by young people. Available at: <https://www.guttmacher.org/state-policy/explore/overview-minors-consent-law>. Accessed August 13, 2018
 64. Alan Guttmacher Institute. State policies on teens. Available at: <https://www.guttmacher.org/united-states/teens/state-policies-teens>. Accessed March 27, 2017
 65. Song X, Klein JD, Yan H, et al. Parent and adolescent attitudes toward preventive care and confidentiality. *J Adolesc Health*. 2019;64(2):235–241
 66. World Health Organization. *Global Standards for Quality Health-Care Services for Adolescents: A Guide to Implement A Standards-Driven Approach to Improve the Quality of Health Care Services for Adolescents*.

- Geneva, Switzerland: World Health Organization; 2015
67. American Academy of Pediatrics, Committee on Medical Liability and Risk Management. In: Donn SM, McAbee GN, eds. *Medicolegal Issues in Pediatrics*, 7th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2005
 68. National Research Council and Institute of Medicine. *A Study of Interactions: Emerging Issues in the Science of Adolescence: Workshop Summary*. Washington, DC: The National Academies Press; 2006
 69. Kessler RC, Berglund P, Demler O, et al. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication [published correction appears in *Arch Gen Psychiatry*. 2005;62(7):768]. *Arch Gen Psychiatry*. 2005;62(6):593–602
 70. Rushton JL, Forcier M, Schectman RM. Epidemiology of depressive symptoms in the National Longitudinal Study of Adolescent Health. *J Am Acad Child Adolesc Psychiatry*. 2002;41(2):199–205
 71. Curtain SC, Heron M, Minino AM, Warner M. Recent increases in injury mortality among children and adolescents aged 10-19 years in the United States: 1999-2016. *Natl Vit Stat Rep*. 2018;67(4):1–16
 72. Centers for Disease Control and Prevention. Trends in the prevalence of suicide-related behaviors. National YRBS: 1991–2017. Available at: https://www.cdc.gov/healthyyouth/data/yrbs/pdf/trends/2017_suicide_trend_yrbs.pdf. Accessed June 26, 2018
 73. Gilreath TD, Wrabel SL, Sullivan KS, et al. Suicidality among military-connected adolescents in California schools. *Eur Child Adolesc Psychiatry*. 2016;25(1):61–66
 74. Rosen DS; American Academy of Pediatrics Committee on Adolescence. Identification and management of eating disorders in children and adolescents. *Pediatrics*. 2010;126(6):1240–1253
 75. Robinson TN, Killen JD, Litt IF, et al. Ethnicity and body dissatisfaction: are Hispanic and Asian girls at increased risk for eating disorders? *J Adolesc Health*. 1996;19(6):384–393
 76. Crago M, Shisslak CM, Estes LS. Eating disturbances among American minority groups: a review. *Int J Eat Disord*. 1996;19(3):239–248
 77. Gard MC, Freeman CP. The dismantling of a myth: a review of eating disorders and socioeconomic status. *Int J Eat Disord*. 1996;20(1):1–12
 78. Kapphahn CJ, Morreale MC, Rickert VI, Walker LR; Society for Adolescent Medicine. Financing mental health services for adolescents: a position paper of the Society for Adolescent Medicine. *J Adolesc Health*. 2006;39(3):456–458
 79. Skowrya KR, Coccozza JJ. *Blueprint for Change: A Comprehensive Model for the Identification and Treatment of Youth with Mental Health Needs in Contact with the Juvenile Justice System*. Delmar, NY: The National Center for Mental Health and Juvenile Justice Policy Research Associates, Inc; 2007
 80. Teplin LA, Abram KM, McClelland GM, Dulcan MK, Mericle AA. Psychiatric disorders in youth in juvenile detention. *Arch Gen Psychiatry*. 2002;59(12):1133–1143
 81. Toro PA, Dworsky A, Fowler PJ. *Toward Understanding Homelessness: The 2007 National Symposium on Homelessness Research*. Washington, DC: US Department of Health and Human Services; 2007
 82. Baird M, Blount A, Brungardt S, et al; Working Party Group on Integrated Behavioral Healthcare. Joint principles: integrating behavioral health care into the patient-centered medical home. *Ann Fam Med*. 2014;12(2):183–185
 83. Lehrer JA, Pantell R, Tebb K, Shafer MA. Forgone health care among U.S. adolescents: associations between risk characteristics and confidentiality concern. *J Adolesc Health*. 2007;40(3):218–226
 84. Kulik DM, Gaetz S, Crowe C, Ford-Jones EL. Homeless youth's overwhelming health burden: a review of the literature. *Paediatr Child Health*. 2011;16(6):e43–e47
 85. Secura GM, Madden T, McNicholas C, et al. Provision of no-cost, long-acting contraception and teenage pregnancy. *N Engl J Med*. 2014;371(14):1316–1323
 86. Baldwin MK, Edelman AB. The effect of long-acting reversible contraception on rapid repeat pregnancy in adolescents: a review. *J Adolesc Health*. 2013;52(suppl 4):S47–S53
 87. Chen XK, Wen SW, Fleming N, et al. Teenage pregnancy and adverse birth outcomes: a large population based retrospective cohort study. *Int J Epidemiol*. 2007;36(2):368–373
 88. Centers for Disease Control and Prevention. *Sexually Transmitted Disease Surveillance 2016*. Atlanta, GA: US Department of Health and Human Services; 2017
 89. McKee MD, Rubin SE, Campos G, O'Sullivan LF. Challenges of providing confidential care to adolescents in urban primary care: clinician perspectives. *Ann Fam Med*. 2011;9(1):37–43
 90. American Academy of Family Physicians, American Academy of Pediatrics, American College of Physicians, American Osteopathic Association. Joint Principles of the patient-centered medical home. 2007. Available at: https://www.aafp.org/dam/AAFP/documents/practice_management/pcmh/initiatives/PCMHJoint.pdf. Accessed March 27, 2017
 91. Cooley WC, Sagerman PJ; American Academy of Pediatrics; American Academy of Family Physicians; American College of Physicians; Transitions Clinical Report Authoring Group. Supporting the health care transition from adolescence to adulthood in the medical home. *Pediatrics*. 2011;128(1):182–200
 92. Committee on Adolescence American Academy of Pediatrics. Achieving quality health services for adolescents. *Pediatrics*. 2008;121(6):1263–1270
 93. Agency for Healthcare Research and Quality. *2006 National*

Healthcare Quality Report. Rockville, MD: US Department of Health and Human Services, Agency for Healthcare Research and Quality; 2006

94. Ginsburg KR, Alexander PM, Hunt J, et al. Enhancing their likelihood for a positive future: the perspective of inner-city youth. *Pediatrics*. 2002;109(6):1136–1142

95. Arenson M, Hudson PJ, Lai B. The evidence on school-based health centers: a review. *Glob Pediatr Health*. 2019;6:2333794X19828745

Unique Needs of the Adolescent

Elizabeth M. Alderman, Cora C. Breuner and COMMITTEE ON ADOLESCENCE
Pediatrics originally published online November 18, 2019;

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/early/2019/11/14/peds.2019-3150
References	This article cites 65 articles, 17 of which you can access for free at: http://pediatrics.aappublications.org/content/early/2019/11/14/peds.2019-3150#BIBL
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Adolescent Health/Medicine http://www.aappublications.org/cgi/collection/adolescent_health_medicine_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://www.aappublications.org/site/misc/reprints.xhtml

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Unique Needs of the Adolescent

Elizabeth M. Alderman, Cora C. Breuner and COMMITTEE ON ADOLESCENCE

Pediatrics originally published online November 18, 2019;

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/early/2019/11/14/peds.2019-3150>

Pediatrics is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. Pediatrics is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2019 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 1073-0397.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

